

# Modena Font Family

NEWLYN

---

Modena has been developed over a period of ten years as a versatile family with clean graphic presence. It's proportions are slightly condensed, having a vertical dynamic that is balanced by the unique horizontal forms of the counters.

The family of five weights and italics has been carefully designed to communicate confidence and authority in a rational and open way.

Modena is suitable for a variety of applications including pharma, finance, transport and culture.

---

Modena has been created by Miles Newlyn, an internationally respected designer, having created identities for some of the world largest organisations, including Honda and Unilever. Newlyn is also an expert in corporate typography, designing typefaces for clients such as Citibank, British Telecom, and the Tate Gallery.



26/28pt

WHEN ON BOARD H.M.S. BEAGLE, AS NATURALIST, I WAS MUCH STRUCK WITH CERTAIN FACTS IN THE DISTRIBUTION

16/18pt

When the first tendency was once displayed, methodical selection and the inherited effects of compulsory training in each successive generation would soon complete the work; and unconscious selection is still at work, as each man tries to procure,

8/10pt

When on board H.M.S. Beagle, as naturalist, I was much struck with certain facts in the distribution of the inhabitants of South America, and in the geological relations of the present to the past inhabitants of that continent. These facts seemed to me to throw some light on the origin of species—that mystery of mysteries, as it has been called by one of our greatest philosophers. As many more individuals of each

species are born than can possibly survive; and as, consequently, there is a frequently recurring struggle for existence, it follows that any being, if it vary however slightly in any manner profitable to itself, under the complex and sometimes varying conditions of life, will have a better chance of surviving, and thus be naturally selected. From the strong principle of inheritance, any selected variety will tend to prop-

6/8pt

Species	C	Common Name	N	1	2	3	4	5	6	7	8	9	10	11
GROUP 01 – DETRITIVORES														
Apereiodon affinis	1	"canivete"	169	0.03	10.16	—	0.01	0.04	0.01	—	17.24	—	72.32	72.32
Apereiodon piracicabae	6	"canivete"	69	0.04	6.66	—	0.05	—	—	—	11.34	1.52	0.74	79.65
Hypostomus ragani	22	"cascudo"	111	—	0.30	0.19	—	—	—	—	3.36	0.01	0.49	95.64
Hypostomus sp.	23	"cascudo"	19	0.01	0.15	—	—	—	—	—	9.50	1.45	—	88.88
Prochilodus lineatus	41	"curimbata"	150	—	0.02	—	0.07	—	—	—	2.05	—	1.58	96.28
Steindachnerina corumbae	46	"saguiru"	289	0.07	0.43	0.04	0.01	—	—	—	3.65	0.01	0.48	95.32
Steindachnerina insculpta	47	"saguiru"	395	0.02	1.22	0.07	0.01	—	—	—	1.93	—	96.62	96.62

26/28pt

*WHEN ON BOARD H.M.S. BEAGLE, AS NATURALIST, I WAS MUCH STRUCK WITH CERTAIN FACTS IN THE DISTRIBUTION OF*

16/18pt

*When the first tendency was once displayed, methodical selection and the inherited effects of compulsory training in each successive generation would soon complete the work; and unconscious selection is still at work, as each man tries to procure, without*

8/10pt

*When on board H.M.S. Beagle, as naturalist, I was much struck with certain facts in the distribution of the inhabitants of South America, and in the geological relations of the present to the past inhabitants of that continent. These facts seemed to me to throw some light on the origin of species—that mystery of mysteries, as it has been called by one of our greatest philosophers. As many more individuals of each species are born than*

*can possibly survive; and as, consequently, there is a frequently recurring struggle for existence, it follows that any being, if it vary however slightly in any manner profitable to itself, under the complex and sometimes varying conditions of life, will have a better chance of surviving, and thus be naturally selected. From the strong principle of inheritance, any selected variety will tend to propagate its new and modified form.*

6/8pt

Species	C	Common Name	N	1	2	3	4	5	6	7	8	9	10	11
GROUP 01 – DETRITIVORES														
<i>Apereiodon affinis</i>	1	"canivete"	169	0.03	10.16	–	0.01	0.04	0.01	–	17.24	–	72.32	72.32
<i>Apereiodon piracicabae</i>	6	"canivete"	69	0.04	6.66	–	0.05	–	–	–	11.34	1.52	0.74	79.65
<i>Hypostomus ragani</i>	22	"cascudo"	111	–	0.30	0.19	–	–	–	–	3.36	0.01	0.49	95.64
<i>Hypostomus sp.</i>	23	"cascudo"	19	0.01	0.15	–	–	–	–	–	9.50	1.45	–	88.88
<i>Prochilodus lineatus</i>	41	"curimbata"	150	–	0.02	–	0.07	–	–	–	2.05	–	1.58	96.28
<i>Steindachnerina corumbae</i>	46	"saguiru"	289	0.07	0.43	0.04	0.01	–	–	–	3.65	0.01	0.48	95.32
<i>Steindachnerina inculpta</i>	47	"saguiru"	395	0.02	1.22	0.07	0.01	–	–	–	1.93	–	96.62	96.62

26/28pt

WHEN ON BOARD H.M.S. BEAGLE, AS NATURALIST, I WAS MUCH STRUCK WITH CERTAIN FACTS IN THE DISTRIBUTION

16/18pt

When the first tendency was once displayed, methodical selection *and the inherited effects of compulsory training in each successive generation* would soon complete the work; and unconscious selection is still at work, as each man tries to pro-

8/10pt

When on board H.M.S. Beagle, as naturalist, I was much struck with certain facts in the distribution of the inhabitants of South America, and in the geological relations of the present to the past inhabitants of that continent. These facts seemed to me to throw some light on the origin of species—that mystery of mysteries, as it has been called by one of our greatest philosophers. As many more individ-

uals of each species are born than can possibly survive; and as, consequently, there is a frequently recurring struggle for existence, it follows that any being, if it vary however slightly in any manner profitable to itself, under the complex and sometimes varying conditions of life, will have a better chance of surviving, and thus be naturally selected. From the strong principle of inheritance, any selected variety will tend

6/8pt

Species	C	Common Name	N	1	2	3	4	5	6	7	8	9	10	11
GROUP 01 – DETRITIVORES														
<i>Apereiodon affinis</i>	1	"canivete"	169	0.03	10.16	–	0.01	0.04	0.01	–	17.24	–	72.32	72.32
<i>Apereiodon piracicabae</i>	6	"canivete"	69	0.04	6.66	–	0.05	–	–	–	11.34	1.52	0.74	79.65
<i>Hypostomus ragani</i>	22	"cascudo"	111	–	0.30	0.19	–	–	–	–	3.36	0.01	0.49	95.64
<i>Hypostomus</i> sp.	23	"cascudo"	19	0.01	0.15	–	–	–	–	–	9.50	1.45	–	88.88
<i>Prochilodus lineatus</i>	41	"curimbata"	150	–	0.02	–	0.07	–	–	–	2.05	–	1.58	96.28
<i>Steindachnerina corumbae</i>	46	"saguiru"	289	0.07	0.43	0.04	0.01	–	–	–	3.65	0.01	0.48	95.32
<i>Steindachnerina inculpta</i>	47	"saguiru"	395	0.02	1.22	0.07	0.01	–	–	–	1.93	–	96.62	96.62

26/28pt

*WHEN ON BOARD H.M.S. BEAGLE, AS NATURALIST, I WAS MUCH STRUCK WITH CERTAIN FACTS IN THE DISTRIBUTION*

16/18pt

*When the first tendency was once displayed, methodical selection and the inherited effects of compulsory training in each successive generation would soon complete the work; and unconscious selection is still at work, as each man tries to procure,*

8/10pt

*When on board H.M.S. Beagle, as naturalist, I was much struck with certain facts in the distribution of the inhabitants of South America, and in the geological relations of the present to the past inhabitants of that continent. These facts seemed to me to throw some light on the origin of species—that mystery of mysteries, as it has been called by one of our greatest philosophers. As many more individuals of each*

*species are born than can possibly survive; and as, consequently, there is a frequently recurring struggle for existence, it follows that any being, if it vary however slightly in any manner profitable to itself, under the complex and sometimes varying conditions of life, will have a better chance of surviving, and thus be naturally selected. From the strong principle of inheritance, any selected variety will tend to propagate*

6/8pt

Species	C	Common Name	N	1	2	3	4	5	6	7	8	9	10	11
GROUP 01 – DETRITIVORES														
<i>Apereiodon affinis</i>	1	"canivete"	169	0.03	10.16	—	0.01	0.04	0.01	—	17.24	—	72.32	72.32
<i>Apereiodon piracicabae</i>	6	"canivete"	69	0.04	6.66	—	0.05	—	—	—	11.34	1.52	0.74	79.65
<i>Hypostomus ragani</i>	22	"cascudo"	111	—	0.30	0.19	—	—	—	—	3.36	0.01	0.49	95.64
<i>Hypostomus sp.</i>	23	"cascudo"	19	0.01	0.15	—	—	—	—	—	9.50	1.45	—	88.88
<i>Prochilodus lineatus</i>	41	"curimbata"	150	—	0.02	—	0.07	—	—	—	2.05	—	1.58	96.28
<i>Steindachnerina corumbae</i>	46	"saguiru"	289	0.07	0.43	0.04	0.01	—	—	—	3.65	0.01	0.48	95.32
<i>Steindachnerina inculpta</i>	47	"saguiru"	395	0.02	1.22	0.07	0.01	—	—	—	1.93	—	96.62	96.62

26/28pt

# WHEN ON BOARD H.M.S. BEAGLE, AS NATURALIST, I WAS MUCH STRUCK WITH CERTAIN FACTS IN THE DISTRIB

16/18pt

When the first tendency was once displayed, methodical selection *and the inherited effects of compulsory training in each successive generation* would soon complete the work; and unconscious selection is still at work, as each man tries

8/10pt

When on board H.M.S. Beagle, as naturalist, I was much struck with certain facts in the distribution of the inhabitants of South America, and in the geological relations of the present to the past inhabitants of that continent. These facts seemed to me to throw some light on the origin of species—that mystery of mysteries, as it has been called by one of our greatest philosophers. As many more

individuals of each species are born than can possibly survive; and as, consequently, there is a frequently recurring struggle for existence, it follows that any being, if it vary however slightly in any manner profitable to itself, under the complex and sometimes varying conditions of life, will have a better chance of surviving, and thus be naturally selected. From the strong principle of inheritance,

6/8pt

Species	C	Common Name	N	1	2	3	4	5	6	7	8	9	10	11
GROUP 01 – DETRITIVORES														
<i>Apereiodon affinis</i>	1	"canivete"	169	0.03	10.16	–	0.01	0.04	0.01	–	17.24	–	72.32	72.32
<i>Apereiodon piracicabae</i>	6	"canivete"	69	0.04	6.66	–	0.05	–	–	–	11.34	1.52	0.74	79.65
<i>Hypostomus ragani</i>	22	"cascudo"	111	–	0.30	0.19	–	–	–	–	3.36	0.01	0.49	95.64
<i>Hypostomus sp.</i>	23	"cascudo"	19	0.01	0.15	–	–	–	–	–	9.50	1.45	–	88.88
<i>Prochilodus lineatus</i>	41	"curimbata"	150	–	0.02	–	0.07	–	–	–	2.05	–	1.58	96.28
<i>Steindachnerina corumbae</i>	46	"saguiru"	289	0.07	0.43	0.04	0.01	–	–	–	3.65	0.01	0.48	95.32
<i>Steindachnerina inculpta</i>	47	"saguiru"	395	0.02	1.22	0.07	0.01	–	–	–	1.93	–	96.62	96.62

26/28pt

**WHEN ON BOARD H.M.S. BEAGLE, AS NATURALIST, I WAS MUCH STRUCK WITH CERTAIN FACTS IN THE DISTRIB**

16/18pt

*When the first tendency was once displayed, methodical selection and the inherited effects of compulsory training in each successive generation would soon complete the work; and unconscious selection is still at work, as each man tries to pro-*

8/10pt

*When on board H.M.S. Beagle, as naturalist, I was much struck with certain facts in the distribution of the inhabitants of South America, and in the geological relations of the present to the past inhabitants of that continent. These facts seemed to me to throw some light on the origin of species—that mystery of mysteries, as it has been called by one of our greatest philosophers. As many more individ-*

*uals of each species are born than can possibly survive; and as, consequently, there is a frequently recurring struggle for existence, it follows that any being, if it vary however slightly in any manner profitable to itself, under the complex and sometimes varying conditions of life, will have a better chance of surviving, and thus be naturally selected. From the strong principle of inheritance, any selected variety will tend*

6/8pt

Species	C	Common Name	N	1	2	3	4	5	6	7	8	9	10	11
GROUP 01 – DETRITIVORES														
<i>Apereiodon affinis</i>	1	"canivete"	169	0.03	10.16	–	0.01	0.04	0.01	–	17.24	–	72.32	72.32
<i>Apereiodon piracicabae</i>	6	"canivete"	69	0.04	6.66	–	0.05	–	–	–	11.34	1.52	0.74	79.65
<i>Hypostomus ragani</i>	22	"cascudo"	111	–	0.30	0.19	–	–	–	–	3.36	0.01	0.49	95.64
<i>Hypostomus sp.</i>	23	"cascudo"	19	0.01	0.15	–	–	–	–	–	9.50	1.45	–	88.88
<i>Prochilodus lineatus</i>	41	"curimbata"	150	–	0.02	–	0.07	–	–	–	2.05	–	1.58	96.28
<i>Steindachnerina corumbae</i>	46	"saguiru"	289	0.07	0.43	0.04	0.01	–	–	–	3.65	0.01	0.48	95.32
<i>Steindachnerina inculpta</i>	47	"saguiru"	395	0.02	1.22	0.07	0.01	–	–	–	1.93	–	96.62	96.62

26/28pt

# WHEN ON BOARD H.M.S. BEAGLE, AS NATURALIST, I WAS MUCH STRUCK WITH CERTAIN FACTS IN THE DISTRIB

16/18pt

When the first tendency was once displayed, methodical selection *and the inherited effects of compulsory training in each successive generation* would soon complete the work; and unconscious selection is still at work, as each man

8/10pt

When on board H.M.S. Beagle, as naturalist, I was much struck with certain facts in the distribution of the inhabitants of South America, and in the geological relations of the present to the past inhabitants of that continent. These facts seemed to me to throw some light on the origin of species—that mystery of mysteries, as it has been called by one of our greatest philosophers. As many

more individuals of each species are born than can possibly survive; and as, consequently, there is a frequently recurring struggle for existence, it follows that any being, if it vary however slightly in any manner profitable to itself, under the complex and sometimes varying conditions of life, will have a better chance of surviving, and thus be naturally selected. From the strong principle

6/8pt

Species	C	Common Name	N	1	2	3	4	5	6	7	8	9	10	11
GROUP 01 – DETRITIVORES														
<i>Apereiodon affinis</i>	1	"canivete"	169	0.03	10.16	–	0.01	0.04	0.01	–	17.24	–	72.32	72.32
<i>Apereiodon piracicabae</i>	6	"canivete"	69	0.04	6.66	–	0.05	–	–	–	11.34	1.52	0.74	79.65
<i>Hypostomus ragani</i>	22	"cascudo"	111	–	0.30	0.19	–	–	–	–	3.36	0.01	0.49	95.64
<i>Hypostomus sp.</i>	23	"cascudo"	19	0.01	0.15	–	–	–	–	–	9.50	1.45	–	88.88
<i>Prochilodus lineatus</i>	41	"curimbata"	150	–	0.02	–	0.07	–	–	–	–	–	1.58	96.28
<i>Steindachnerina corumbae</i>	46	"saguiru"	289	0.07	0.43	0.04	0.01	–	–	–	3.65	0.01	0.48	95.32
<i>Steindachnerina insculpta</i>	47	"saguiru"	395	0.02	1.22	0.07	0.01	–	–	–	1.93	–	96.62	96.62

26/28pt

**WHEN ON BOARD H.M.S. BEAGLE, AS NATURALIST, I WAS MUCH STRUCK WITH CERTAIN FACTS IN THE DISTRIB**

16/18pt

***When the first tendency was once displayed, methodical selection and the inherited effects of compulsory training in each successive generation would soon complete the work; and unconscious selection is still at work, as each man tries***

8/10pt

*When on board H.M.S. Beagle, as naturalist, I was much struck with certain facts in the distribution of the inhabitants of South America, and in the geological relations of the present to the past inhabitants of that continent. These facts seemed to me to throw some light on the origin of species—that mystery of mysteries, as it has been called by one of our greatest philosophers. As many more*

*individuals of each species are born than can possibly survive; and as, consequently, there is a frequently recurring struggle for existence, it follows that any being, if it vary however slightly in any manner profitable to itself, under the complex and sometimes varying conditions of life, will have a better chance of surviving, and thus be naturally selected. From the strong principle of inherit-*

6/8pt

Species	C	Common Name	N	1	2	3	4	5	6	7	8	9	10	11
GROUP 01 – DETRITIVORES														
<i>Apereiodon affinis</i>	1	"canivete"	169	0.03	10.16	–	0.01	0.04	0.01	–	17.24	–	72.32	72.32
<i>Apereiodon piracicabae</i>	6	"canivete"	69	0.04	6.66	–	0.05	–	–	–	11.34	1.52	0.74	79.65
<i>Hypostomus ragani</i>	22	"cascudo"	111	–	0.30	0.19	–	–	–	–	3.36	0.01	0.49	95.64
<i>Hypostomus sp.</i>	23	"cascudo"	19	0.01	0.15	–	–	–	–	–	9.50	1.45	–	88.88
<i>Prochilodus lineatus</i>	41	"curimbata"	150	–	0.02	–	0.07	–	–	–	–	–	1.58	96.28
<i>Steindachnerina corumbae</i>	46	"saguiru"	289	0.07	0.43	0.04	0.01	–	–	–	3.65	0.01	0.48	95.32
<i>Steindachnerina insculpta</i>	47	"saguiru"	395	0.02	1.22	0.07	0.01	–	–	–	1.93	–	96.62	96.62

26/28pt

# WHEN ON BOARD H.M.S. BEAGLE, AS NATURALIST, I WAS MUCH STRUCK WITH CERTAIN FACTS IN

16/18pt

When the first tendency was once displayed, methodical selection *and the inherited effects of compulsory training in each successive generation* would soon complete the work; and unconscious selection is still at work, as

8/10pt

When on board H.M.S. Beagle, as naturalist, I was much struck with certain facts in the distribution of the inhabitants of South America, and in the geological relations of the present to the past inhabitants of that continent. These facts seemed to me to throw some light on the origin of species—that mystery of mysteries, as it has been called by one of our greatest phi-

losophers. As many more individuals of each species are born than can possibly survive; and as, consequently, there is a frequently recurring struggle for existence, it follows that any being, if it vary however slightly in any manner profitable to itself, under the complex and sometimes varying conditions of life, will have a better chance of surviving, and thus be natu-

6/8pt

Species	C	Common Name	N	1	2	3	4	5	6	7	8	9	10	11
GROUP 01 – DETRITIVORES														
<i>Apereiodon affinis</i>	1	"canivete"	169	0.03	10.16	–	0.01	0.04	0.01	–	17.24	–	72.32	72.32
<i>Apereiodon piracicabae</i>	6	"canivete"	69	0.04	6.66	–	0.05	–	–	–	11.34	1.52	0.74	79.65
<i>Hypostomus ragani</i>	22	"cascudo"	111	–	0.30	0.19	–	–	–	–	3.36	0.01	0.49	95.64
<i>Hypostomus sp.</i>	23	"cascudo"	19	0.01	0.15	–	–	–	–	–	9.50	1.45	–	88.88
<i>Prochilodus lineatus</i>	41	"curimbata"	150	–	0.02	–	0.07	–	–	–	2.05	–	1.58	96.28
<i>Steindachnerina corumbae</i>	46	"saguiru"	289	0.07	0.43	0.04	0.01	–	–	–	3.65	0.01	0.48	95.32
<i>Steindachnerina insculpta</i>	47	"saguiru"	395	0.02	1.22	0.07	0.01	–	–	–	1.93	–	96.62	96.62

26/28pt

**WHEN ON BOARD H.M.S. BEAGLE, AS NATURALIST, I WAS MUCH STRUCK WITH CERTAIN FACTS IN THE DISTRIB**

16/18pt

**When the first tendency was once displayed, methodical selection and the inherited effects of compulsory training in each successive generation would soon complete the work; and unconscious selection is still at work, as each**

8/10pt

*When on board H.M.S. Beagle, as naturalist, I was much struck with certain facts in the distribution of the inhabitants of South America, and in the geological relations of the present to the past inhabitants of that continent. These facts seemed to me to throw some light on the origin of species—that mystery of mysteries, as it has been called by one of our greatest phi-*

*losophers. As many more individuals of each species are born than can possibly survive; and as, consequently, there is a frequently recurring struggle for existence, it follows that any being, if it vary however slightly in any manner profitable to itself, under the complex and sometimes varying conditions of life, will have a better chance of surviving, and thus be naturally selected.*

6/8pt

Species	C	Common Name	N	1	2	3	4	5	6	7	8	9	10	11
<b>GROUP 01 – DETRITIVORES</b>														
<i>Apereiodon affinis</i>	1	"canivete"	169	0.03	10.16	–	0.01	0.04	0.01	–	17.24	–	72.32	72.32
<i>Apereiodon piracicabae</i>	6	"canivete"	69	0.04	6.66	–	0.05	–	–	–	11.34	1.52	0.74	79.65
<i>Hypostomus ragani</i>	22	"cascudo"	111	–	0.30	0.19	–	–	–	–	3.36	0.01	0.49	95.64
<i>Hypostomus sp.</i>	23	"cascudo"	19	0.01	0.15	–	–	–	–	–	9.50	1.45	–	88.88
<i>Prochilodus lineatus</i>	41	"curimbata"	150	–	0.02	–	0.07	–	–	–	2.05	–	1.58	96.28
<i>Steindachnerina corumbae</i>	46	"saguiru"	289	0.07	0.43	0.04	0.01	–	–	–	3.65	0.01	0.48	95.32
<i>Steindachnerina insculpta</i>	47	"saguiru"	395	0.02	1.22	0.07	0.01	–	–	–	1.93	–	96.62	96.62